

University of Hawai'i at Mānoa LIBRARIES



Interlibrary Loan Program

Hamilton Library, Room 101 ♦ 2550 McCarthy Mall ♦ Honolulu, HI 96822

Phone/Voicemail: 808-956-5956 (Lending) / 808-956-8568 (for UHM patrons)

Fax: 808-956-5968

Email: lendill@hawaii.edu (Lending) / libill@hawaii.edu (for UHM patrons)



NOTICE: The copyright law of the United States (Title 17, U.S.C.) governs the making of photocopies or other reproductions of copyrighted materials. Under certain conditions specified in the law, libraries and archives are authorized to furnish a photocopy or reproduction. One of these specified conditions is that the photocopy or reproduction is not to be “used for any purpose other than in private study, scholarship, or research.” If a user makes a request for, or later uses, a photocopy or reproduction for purposes in excess of “fair use,” that user may be liable for copyright infringement. This institution reserves the right to refuse to accept a copying order, if, in its judgment, fulfillment of the order would involve violation of copyright law.



BEST AVAILABLE COPY

University of Hawai'i at Mānoa LIBRARIES

Interlibrary Loan Program

Hamilton Library Room 101 ♦ 2550 McCarthy Mall ♦ Honolulu, HI 96822
Phone 461-2200 ♦ Fax 461-2201 ♦ E-mail lib@hawaii.gov

Phone/Voicemail: 808-956-5956

Fax: 808-956-5968

Email: lendill@hawaii.edu

NOTICE: The copyright law of the United States (Title 17, U.S.C.) governs the making of photocopies or other reproductions of copyrighted materials. Under certain conditions specified in the law, libraries and archives are authorized to furnish a photocopy or reproduction. One of these specified conditions is that the photocopy or reproduction is not to be "used for any purpose other than in private study, scholarship, or research." If a user makes a request for, or later uses, a photocopy or reproduction for purposes in excess of "fair use," that user may be liable for copyright infringement. This institution reserves the right to refuse to accept a copying order, if, in its judgment, fulfillment of the order would involve violation of copyright law.

BEST-AVAILABLE COPY

UNIVERSITY OF HAWAII
LIBRARY

The Chemical Society, London

Chemical Communications

6
1968

ICAL

DI

88

apart the *peri*-CH₂ groups, and consequently distort the aromatic system. The strain involved causes the measurable barrier to nitrogen inversion, and this novel demonstration of the *peri*-effect.

¹ V. Balasubramaniyam, *Chem. Rev.*, 1966, **66**, 567.

² G. Wittig and H. Ludwig, *Annalen*, 1954, **589**, 55.

³ L. W. Reeves, *Adv. Phys. Org. Chem.*, 1965, **3**, 187.

⁴ D. L. Griffith and J. D. Roberts, *J. Amer. Chem. Soc.*, 1965, **87**, 4089; J. E. Anderson and J. M. Lehn, *ibid.*, 1967, **89**, 81.

⁵ J. M. Lehn, F. G. Riddell, B. J. Price, and I. O. Sutherland, *J. Chem. Soc. (B)*, 1967, 387.

⁶ J. Loewenstein, J. F. Neumer, and J. D. Roberts, *J. Amer. Chem. Soc.*, 1960, **82**, 3599.

This work was supported by the National Science Foundation, and the National Research Council, Canada.

(Received, January 26th, 1968; Com. 098.)

The Thermal Decomposition of 1,3,5-Trinitrohexahydro-1,3,5-triazine (RDX)

By J. D. COSGROVE and A. J. OWEN*†

(Explosives Research and Development Establishment, Ministry of Technology, Waltham Abbey, Essex)


THE pressure-time curve obtained for the thermal decomposition of RDX below its melting point (204°) has been interpreted¹ as a slow solid-state reaction leading to a product which liquifies the reactant at the temperature of the experiment and accelerates its decomposition. The ratio of the rates of decomposition in the liquid and solid phases is said to be approximately 10 : 1. We have studied the decomposition and conclude that the

initial reaction occurs in the gas phase and not in the solid state.

Reaction flasks² of volumes between 11 and 150 ml. containing 0.02–0.2 g. RDX were either evacuated or filled to a known pressure with nitrogen and sealed. They were completely immersed in a bath at 195° for 1 hr., cooled to room temperature, opened under vacuum to determine the pressure of gaseous products, and the residual RDX estimated by nonaqueous titration.³ The results, illustrated in the Figure, show that the rate of decomposition is (a) directly proportional to the volume of the reaction vessel; (b) for a constant volume, independent of the amount of RDX present; and (c) retarded by the presence of inert gases *e.g.* nitrogen.

It is clear from (a) and (b) that RDX does not decompose in the solid state to any significant extent at this temperature. In the early stages of the decomposition, RDX exerts its equilibrium vapour pressure⁴ and the gas then decomposes into several products including a liquid, hydroxymethylformamide which dissolves some of the RDX and results in an accelerating reaction. The rate of reaction in the gas phase is at least equal to that in the liquid phase, but since the overall rate is controlled by the vapour pressure of RDX, and its value is not known with any degree of accuracy at present, an absolute value for the rate constant for the gas-phase reaction cannot be calculated.

(Received, January 1st, 1968; Com. 006.)



FIGURE

Initial conditions: + 0.200 g. RDX, evacuated; ● 0.100 g. RDX, evacuated; △ 0.072 g. RDX, evacuated; ○ 0.020 g. RDX, evacuated; □ 0.200 g. RDX, 58 mm. nitrogen; ■ 0.200 g. RDX, 119 mm. nitrogen.

* Present address, Royal Armament Research and Development Establishment, Ministry of Defence, Woolwich, London, S.E.18.

¹ C. E. H. Bawn, "Chemistry of the Solid State," Butterworths, London, 1955, pp. 261, 263.

² A. J. Owen, *J. Chem. Soc.*, 1961, 5438.

³ S. M. Kaye, *Analyt. Chem.*, 1955, **27**, 292.

⁴ G. Edwards, *Trans. Faraday Soc.*, 1953, **49**, 152.