

AD-752 899

INFRARED AND RAMAN SPECTRA OF 1, 3, 5-  
TRINITRO-1, 3, 5-TRIAZACYCLOHEXANE (RDX)

Zafar Iqbal, et al

Picatinny Arsenal  
Dover, New Jersey

October 1972

DISTRIBUTED BY:

**NTIS**

National Technical Information Service  
U. S. DEPARTMENT OF COMMERCE  
5285 Port Royal Road, Springfield Va. 22151

AD752899

COPY NO. 21

TECHNICAL REPORT 4401

INFRARED AND RAMAN SPECTRA  
OF

1, 3, 5-TRINITRO-1, 3,  
5-TRIAZACYCLOHEXANE (RDX)

ZAFAR IQBAL  
K. SURYANARAYANAN  
SURYANARAYANA EILUSU  
J. R. AUTERA


OCTOBER 1972



APPROVED FOR PUBLIC RELEASE; DISTRIBUTION UNLIMITED.

Reproduced by  
NATIONAL TECHNICAL  
INFORMATION SERVICE  
U.S. Department of Commerce  
Springfield, VA 22151

PICATINNY ARSENAL  
DOVER, NEW JERSEY




128

The findings in this report are not to be construed  
as an official Department of the Army position.

#### DISPOSITION

Destroy this report when it is no longer needed Do  
not return it to the originator.



**UNCLASSIFIED**  
Security Classification

DOCUMENT CONTROL DATA - R & D		
<i>(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)</i>		
1. ORIGINATING ACTIVITY (Corporate author)		2a. REPORT SECURITY CLASSIFICATION
U.S. Army, Picatinny Arsenal, Dover, NJ		Unclassified
		2b. GROUP
3. REPORT TITLE		
Infrared and Raman Spectra of 1,3,5-Trinitro-1,3,5-Triazacyclohexane (RDX)		
4. DESCRIPTIVE NOTES (Type of report and inclusive dates)		
5. AUTHOR(S) (First name, middle initial, last name)		
Zafar Iqbal; K. Suryanarayana; Suryanarayana Bulusu; J.R. Autera		
6. REPORT DATE	7a. TOTAL NO. OF PAGES	7b. NO. OF REFS
8a. CONTRACT OR GRANT NO.		8b. ORIGINATOR'S REPORT NUMBER(S)
AMCMS Code 501B.11.85601		Technical Report 4401
a. PROJECT NO.		8c. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)
c. DA Project 1T061102A32B 01		
d.		
10. DISTRIBUTION STATEMENT		
Approved for public release; distribution unlimited.		
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY
		Picatinny Arsenal, Dover, NJ
13. ABSTRACT		
<p>The transmission infrared and the laser excited Raman spectra of polycrystalline RDX have been measured in the range of 40 to 4000 <math>\text{cm}^{-1}</math>. To aid assignments in the spectral region of 400 to 4000 <math>\text{cm}^{-1}</math>, the spectra of two types of <math>\text{N}^{15}</math> - labeled samples and the solution spectra in different solvents were also recorded. From these data it was possible to assign many of the observed bands to intramolecular modes of the RDX molecule. The Raman active lattice modes also were resolved and found to be comparable to the lattice mode frequencies in solid cyclohexane.</p>		

DD FORM 1473  
1 NOV 66

REPLACES DD FORM 1473, 1 JAN 64, WHICH IS  
OBSOLETE FOR ARMY USE.

**UNCLASSIFIED**  
Security Classification

I

UNCLASSIFIED

Security Classification

14. KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
Infrared spectra Raman spectra Vibrational spectra 1,3,5-trinitro-1,3,5-triazacyclohexane (RDX) CH <sub>2</sub> modes NO <sub>2</sub> modes Ring vibrations Lattice modes Combination and overtone modes N <sup>15</sup> - labeled RDX Isotopic shift 1,3,5-trinitro-1,3,5-triazacyclohexane (TNTC)						

UNCLASSIFIED

Security Classification

11

Technical Report 4401

INFRARED AND RAMAN SPECTRA OF  
1,3,5-TRINITRO-1,3,5-TRIAZACYCLOHEXANE (RDX)

by

Zafar Iqbal  
K. Suryanarayanan  
Suryanarayana Bulusu  
J.R. Autera

October 1972

AMCMS Code 501B.11.85601  
DA Project 1T061102A32B 01

Approved for public release; distribution  
unlimited.

Explosives Division  
Feltman Research Laboratory  
Picatinny Arsenal  
Dover, New Jersey

III

The citation in this report of the trade names of commercially available products does not constitute official indorsement or approval of the use of such products.

## TABLE OF CONTENTS

	Page No.
Abstract	1
Introduction	2
Symmetry Considerations and Structure	2
Experimental	5
Results	6
Discussion	7
Fundamental Molecular Vibrations	7
Lattice Modes	9
Combination and Overtone Modes	9
Conclusion	9
References	10
Distribution List	21
Tables	
1    Frequencies and assignments of bands in the 40 to 4000 $\text{cm}^{-1}$ region in the spectrum of RDX	11
2    Combination and overtone bands observed in the 1700-3000 $\text{cm}^{-1}$ region in the trans- mission infrared spectrum of a thick single crystal of RDX	14



## Figures

1	Structure of the molecule at a crystal site	16
2	% Transmission (T) vs frequency traces of the infrared spectra of RDX in the 400 to 4000 $\text{cm}^{-1}$ region	17
3	Traces of the Raman spectra of RDX in the 400 to 4000 $\text{cm}^{-1}$ region	18
4	% Transmission (T) vs frequency traces of the KBr pellet infrared spectra between 400 and 1700 $\text{cm}^{-1}$	19
5	Traces of the low frequency infrared and Raman spectra	20

## ABSTRACT

The transmission infrared and the laser excited Raman spectra of polycrystalline RDX have been measured in the range of 40 to 4000  $\text{cm}^{-1}$ . To aid assignments in the spectral region of 400 to 4000  $\text{cm}^{-1}$ , the spectra of two types of  $\text{N}^{15}$  - labeled samples and the solution spectra in different solvents were also recorded. From these data it was possible to assign many of the observed bands to intramolecular modes of the RDX molecule. The Raman active lattice modes also were resolved and found to be comparable to the lattice mode frequencies in solid cyclohexane.

## INTRODUCTION

The nitramines form an interesting group of organic compounds, which have provided structural chemists with a wide variety of problems. This report deals with RDX (1,3,5-trinitro-1,3,5-triazacyclohexane), a monocyclic member of the above group, consisting of three nitramine units. The molecule has 21 atoms; it was thought that it would show sufficient symmetry to allow an analysis of its vibrational spectrum. Such an analysis would permit qualitative conclusions regarding molecular bond orders and conformations in different environments. Previous studies include a detailed structural analysis of the compound by X-ray diffraction (Ref 1) as well as by neutron scattering (Ref 2), and a partial analysis of the mid-infrared part of the spectrum (Ref 3,4). While the present work has been in progress, another study of the infrared and Raman spectra of RDX has been published by Rey-Lafon (Ref 5) et al. Their data are in general agreement with ours; the purpose of the concise account presented here is only to stress the more unusual aspects of the RDX structure. The objective of this work has been to measure the infrared and Raman active vibrations in as wide a spectral range as feasible, and assign the modes by analogy with previous studies on similar but simpler molecules, and from isotopic and solution data. The results obtained may, therefore, form a starting point for a normal coordinate analysis of the molecular spectrum of RDX and the vibrational analysis of the related eight-membered ring HMX molecule (1,3,5,7-tetranitro-1,3,5,7-tetrazacyclooctane).

## SYMMETRY CONSIDERATIONS AND STRUCTURE

RDX has a structure similar to the boat and chair forms of cyclohexane, and as in the latter the chair form should be of lower energy. It is, therefore, expected that RDX will exist predominantly in the chair form. The structure as determined by X-ray diffraction (Ref 1) and confirmed by neutron scattering (Ref 2) of crystalline RDX is shown in Fig 1. This does represent a chair form but, surprisingly, with an unfavorable orientation of two nitramine groups in the axial positions.

In solution, the symmetry of the molecule is, due to ring inversion, that of a symmetric top belonging to point symmetry  $C_{3v}$ . If the N and O atoms of the nitramine group in position 1 (see Fig 1) are in the same plane as the hydrogen atoms of the C atom in position 4, then the 57 normal mode vibrations of the molecule are distributed among the following symmetry species:

$$\Gamma^I = 13 A_1 + 6 A_2 + 19 E$$

However, if the O atoms of the N-NO<sub>2</sub> group in position 1 are not in that plane, the vibrations are distributed as:

$$\Gamma^{II} = 12 A_1 + 7 A_2 + 19 E$$

The vibrations of  $A_1$  and E symmetry are active in both the infrared and Raman spectra, while vibrations of  $A_2$  symmetry are silent in both. Thus, in structure I (symmetry species  $\Gamma^I$ ), 32 modes, in structure II (symmetry species  $\Gamma^{II}$ ), 31 modes are active. The comparatively short N-N bond length of the equatorial N-NO<sub>2</sub> group observed in the crystal (Ref 1) would favor structure II in the solid phase. This may also be true for the isolated molecule.

In the condensed phase RDX loses its threefold symmetry. A staggered chair conformation as shown in Fig 1, in which 2 axial and 1 equatorial -NO<sub>2</sub> groups are found, is suggested by the X-ray diffraction data (Ref 1). The molecules are at sites of symmetry  $C_1$  in the space group  $D_{2h}^{15}$ -Pbca with eight formula units in the primitive cell. According to the site group approximation, the  $A_2$  type molecular vibrations will, therefore, have both infrared and Raman active components in the solid state spectra. The external or lattice mode optic translatory (T) and libratory (R) components for the factor group  $D_{2h}^{15}$ , will be distributed as:

$$\begin{aligned} \Gamma = & (3T + 3R) A_{1g} + (3T + 3R) B_{1g} + (3T + 3R) B_{2g} + \\ & (3T + 3R) B_{3g} + (3T + 3R) A_{1u} + (2T + 3R) B_{1u} + \\ & (2T + 3R) B_{2u} + (2T + 3R) B_{3u} \end{aligned}$$

The factor group is centrosymmetric; hence the Raman active vibrations would be forbidden in the infrared. In principle, however, the correlation field in organic crystals is small, and hence the infrared and Raman spectra in the condensed phase would be nearly coincident. The large number of factor group splittings predicted by group theory would not be observed, either, since the splittings predicted may amount to less than  $1 \text{ cm}^{-1}$ .

## EXPERIMENTAL

The infrared spectra in the region 400 to 4000  $\text{cm}^{-1}$  were recorded in a Perkin-Elmer 621 grating spectrometer using KBr disc samples and solutions. Single crystal measurements were performed using the spectrometer with 6x beam condensing arrangement. The infrared spectra of the region 70 to 400  $\text{cm}^{-1}$  were measured in a Beckman IR-11 spectrometer, using a thin film of the sample in Nujol oil on a polyethylene substrate.

The Raman spectra were recorded using both He-Ne (6328 Å, 50 mW) and Ar ion (4830 Å, 300 nW) focussed radiation. The 90° scattered radiation was analyzed by a Spex 1400 double monochromator coupled with a photon counting detection system. The powder and solution spectra were measured in glass capillaries.

The RDX samples used were purified by multiple recrystallization. Single crystals were grown from acetone solutions of the compound. The ring  $\text{N}^{15}$ -labeled RDX samples were prepared by the nitration of  $\text{N}^{15}$ -labeled hexamine, while the nitro-group labeled RDX samples were prepared by nitrating hexamine with  $\text{N}^{15}$ -labeled nitric acid. The isotopic enrichment in all samples was about 99%.

## RESULTS

The infrared and Raman spectra of RDX as single crystal, powder, and solution samples, have been measured in the frequency range 400 to 4000  $\text{cm}^{-1}$ . The tracings of the spectra are shown in Figures 2 and 3. The solution Raman spectra were recorded using saturated solutions of RDX in deuterated dimethylsulphoxide ( $\text{d}_6$ -DMSO), while the solution infrared spectra in the frequency ranges 1200 to 3200  $\text{cm}^{-1}$  and 600 to 1200  $\text{cm}^{-1}$ , were recorded using as solvents  $\text{d}_6$ -DMSO and acetonitrile, respectively. The powder infrared spectra of ring and nitro group  $\text{N}^{15}$  labeled RDX samples, in the spectral range 400 to 1800  $\text{cm}^{-1}$ , also were measured; the tracings are shown in Fig 4 together with the spectrum for unlabeled RDX. The low frequency 40 to 400  $\text{cm}^{-1}$  region of the spectrum was recorded only for solid samples of RDX and is shown in Fig 5. The infrared spectrum of 1,3,5-trinitroso-1,3,5-triazacyclohexane (TNTC), the trinitroso analog of RDX also was measured; its spectrum in the low frequency region is shown in Fig 5b.

The frequencies of the solid and solution bands of RDX together with the assignments are listed in Table 1. Since a wider frequency range has been measured with powder samples, the frequencies quoted in Table 1 have been taken from the powder spectra. The combination and overtone frequencies in the spectral region 1700 to 3000  $\text{cm}^{-1}$  were observed in the transmission infrared spectrum of a thick single crystal. These frequencies have been compiled in Table 2.

## DISCUSSION

### Fundamental Molecular Vibrations

#### CH<sub>2</sub> Modes

The bands in the 2900 to 3100  $\text{cm}^{-1}$  region can be assigned to CH<sub>2</sub> stretching modes. An assignment in terms of symmetric and asymmetric motions is based on the fact that the symmetric mode gives rise to strong bands in the Raman, and weaker peaks in the infrared spectrum. A number of interesting features can be noted on comparing the solution with the solid infrared and Raman spectra. The higher frequency asymmetric stretching mode in the solid shows a site group splitting of about 10  $\text{cm}^{-1}$  (Fig 2d). This is in agreement with the presence of two types of sites in the staggered chair conformation of the molecule in the crystal. Two peaks separated by 50  $\text{cm}^{-1}$  are observed in the solid state spectra in the region of the symmetric stretching mode. Since the second peak disappears in the spectra of RDX in d<sub>6</sub>-DMSO solution, one can associate this band with the lowered symmetry of the molecule at the crystal site, which leads to the infrared and Raman activity of this mode. In d<sub>6</sub>-DMSO solution, however, both the symmetric and asymmetric stretching peaks broaden out, indicating the presence of possible solvent-solute interaction in that solution.

A point worth noting is that the mode frequency for the CH<sub>2</sub> asymmetric stretching motion in RDX is about 100  $\text{cm}^{-1}$  higher than in cyclohexane. This anomalous value seems to indicate considerable strain on the RDX ring in the crystalline state. This aspect of RDX structure is the subject of a more detailed study involving a group of cyclic nitramines, which will be published shortly.

The assignment of the band between 1390 and 1460  $\text{cm}^{-1}$  to CH<sub>2</sub> skeletal motions is based on the data of Wood (Ref 3) on deuterated RDX.



### NO<sub>2</sub> Modes

The symmetric and asymmetric stretching modes of the NO<sub>2</sub> group can be assigned in the same manner as the corresponding motions in the CH<sub>2</sub> group. These bands would also show the largest shift when the nitrogen atom on the NO<sub>2</sub> group is labeled with N<sup>15</sup>. The peaks between 1540 to 1600 cm<sup>-1</sup> can be clearly identified as the asymmetric NO<sub>2</sub> stretching mode components. The extremely strong peak at approximately 1596 cm<sup>-1</sup> shows a splitting of between 10 and 15 cm<sup>-1</sup>, which can be attributed to the two types of NO<sub>2</sub> groups in the staggered chair conformation of the molecule in the crystal.

The symmetric-stretching NO<sub>2</sub> and the N-N vibrations can have mode frequencies close to one another (Ref 6). Since the peak at 1268 cm<sup>-1</sup> is very strong in the solution Raman spectrum, one can probably assign this band to the NO<sub>2</sub> type stretching. The peak at 1310 cm<sup>-1</sup> can, therefore, be associated with the symmetric stretching N-N mode. Both these modes show a site group splitting of approximately 10 cm<sup>-1</sup> in the infrared absorption measurements on a solid sample.

Two bands in the 750 cm<sup>-1</sup> region of the spectrum show a large shift for nitro group labeled RDX (see Fig 4) and can therefore be assigned to a NO<sub>2</sub> vibration, possibly the NO<sub>2</sub> deformation motion. The skeletal NO<sub>2</sub> modes lie in the low frequency region of the spectrum and are probably mixed to a large degree with torsional and bending ring vibrations. Hence, assignments on the basis of frequency shifts of the nitro group labeled RDX samples would be difficult. However, by comparison with the spectrum of TNTC one can assign the bands between 450 to 610 cm<sup>-1</sup>, which do not occur in the spectrum of the trinitroso analog (see Fig 5b), to NO<sub>2</sub> skeletal motions. The broad bands between 128 and 150 cm<sup>-1</sup> are somewhat too high in energy to be associated with lattice vibrations (see the subsection on lattice modes) and hence can be tentatively identified as due to the N-NO<sub>2</sub> torsional mode.

### Ring Vibrations

The strongest band in the Raman spectrum of solid RDX occurs at  $832\text{ cm}^{-1}$ . This peak can be assigned to the symmetric ring stretching mode since such a motion is likely to generate a large polarizability change. By analogy with the work on cyclohexane (Ref 5), the bands around  $200\text{ cm}^{-1}$  can be assigned to a ring bending and torsional motion, while the peak at  $410\text{ cm}^{-1}$ , which corresponds closely to the strongest band in the neutron inelastic scattering (NIS) spectrum of the salt, can be assigned to an essentially ring bending vibration.

Two sets of frequencies at  $1020$  and  $1220\text{ cm}^{-1}$  have been assigned to ring type vibrations on the basis of the observed shifts for the ring labeled RDX sample.

### Lattice Modes

Five peaks have been resolved in the solid state Raman spectrum between  $50$  and  $110\text{ cm}^{-1}$ , which can be assigned to the external optic or lattice modes. These peaks correspond to the broad density of states centered at  $55\text{ cm}^{-1}$  (Ref 8).

### Combination and Overtone Modes

The bands between  $1700$  and  $3000\text{ cm}^{-1}$ , observed in the infrared spectrum of a thick single crystal of RDX, can be assigned to combination and overtone modes. The spectrum (see Fig 2b) is divided into four regions, as shown in the figure. Possible overtone and summation modes can be suggested for these bands in terms of the strongest infrared and Raman peaks. These are shown in Table 2.

## CONCLUSION

Possible assignments of the rich infrared and Raman spectra of RDX, both in the solid and in  $d_6$ -DMSO and acetonitrile solutions, have been proposed. Site group splittings, observed in the solid state spectra for the  $\text{CH}_2$  asymmetric stretching and the  $\text{NO}_2$  and N-N stretching modes, support the proposed staggered chair conformation of the molecule in the crystal with two distinctive

orientations of the nitramine groups. However, solution data suggest a more symmetrical RDX conformation than in the crystal lattice and some possible solute-solvent interaction. A complete normal coordinate analysis of RDX molecule is being carried out in collaboration with Dr. George Adams of the Propellants Division, Feltman Research Laboratory. A detailed structural study of HMX polymorphs by infrared and Raman spectra also is in progress.

#### REFERENCES

1. P. M. Harris, "Crystal Structures of Aromatic Trinitro and Related Compounds", AFOSR-TR-59-165, Ch 2, September 1959
2. Chang S. Choi, Acta Cryst., in print
3. R. Wood, Proc. Intern. Meet. Mol. Spectroscopy, (4th meeting), Bologna, 2, 955 (1959)
4. D. E. Davey and J. Eadie, Tech. Note. No. 129, Australian Defence Scientific Service, August 1969
5. Madeleine Rey-Lafon, et al, J. de Chim. Phys., 68, 1533, 1573 (1971)
6. V. G. Avakyan and V.A. Chekrygin, Izv. Akad. Nauk SSR, Ser. Khim., (10) 2194 (1969)
7. H. Takahashi and T. Shimanouchi, J. Mol. Spectro., 13, 43, (1964)
8. H. Boutin, S. Trevino and H. Prask, Picatinny Arsenal Technical Report 3370, (1966)

TABLE I  
Frequencies and assignments of bands  
in the 40 to 4000  $\text{cm}^{-1}$  region in the spectrum of RDX

<u>Infrared(<math>\text{cm}^{-1}</math>)</u>		<u>Raman(<math>\text{cm}^{-1}</math>)</u>		<u>Remarks and Assignments</u>
<u>Solid</u>	<u>Solution</u>	<u>Solid</u>	<u>Solution</u>	
		50.3 s <sup>a</sup>		Lattice modes
		59.2 vs		
		71.5 s		
		85.7 m		
		107 m		N-NO <sub>2</sub> torsion
		128 m br		
		148 m br		
		183 w		
225 ms		206 m		C-N-C bend + torsion
		224 s		
346 s		349 s		
412 m		410 w	410 m	C-N-C bend
460 m		464 m	470 w	NO <sub>2</sub> skeletal
487 m		487 w br	490 w	
595 m	595 m	590 m <sup>b</sup>	sol. <sup>c</sup>	
610 m	610 m sh	605 m <sup>b</sup>	sol.	
675 w		671 w	sol.	
740 w sh	740 w	738 w	sol.	NO <sub>2</sub> deformation
754 m	750 m	758 w	sol.	

TABLE 1 (cont'd)

782 m	794 m	787 w	790 m	
844 m		847 m	845 m	
854 m		854 w sh	850 m sh	
882 m	885 w sh	880 vs	885 ms	Symmetric ring stretch
917 vs				
927 vs	910 m	916 w		
947 s	935 s	940 m	947 w	
1020 m	1015 s			
1040 s	1045 m sh	1030 m	sol.	Ring
1219 m		1217 s	1214 m	Ring
1232 m	1230 m sh	1232 w sh		
1267 s				
1275 s sh	1270 s	1274 s	1268 vs	NO <sub>2</sub> symmetric stretch
1312 m				
1322 m	1320 m	1313 s	1309 m	N-N symmetric stretch
1352 m		1344 w	1339 w	
1391 m	1392 m	1382 m sh	1384 m	CH <sub>2</sub> skeletal modes
		1389 m		
1424 m				
	1435 m	1427 m sh	1434 m	
1435 m				
1460 m	1460 m	1461 w		
		1509 w		
1535 ms	1550 m sh	1544 m	1542 m sh	NO <sub>2</sub> asymmetric stretch
		1574 m		
1576 s		1577 m sh		
1593 s	1585 s	1596 m	1582 s	

TABLE 1 (cont'd)

2950 w <sup>b</sup>		2948 vs		] CH <sub>2</sub> symmetric stretch
	2980 w		2967 s	
3006 m		3004 s		] CH <sub>2</sub> asymmetric stretch
3068 ms		3067 m		
3078 s	3080 m	3074 m	3074 m	

<sup>a</sup> s-strong, m-medium strong, w-weak, sh-shoulder, br-broad

<sup>b</sup> observed in single crystal sample only

<sup>c</sup> solvent interference

TABLE 2

Combination and Overtone bands observed  
in the 1700-3000  $\text{cm}^{-1}$  region in the transmission  
infrared spectrum of a thick single crystal of RDX

<u>Frequency (<math>\text{cm}^{-1}</math>)</u>	<u>Remarks</u>
1770 ms	2 x 885
1875 ms	1592 + 1267
1915 m br	Internal-external combination peaks; 1875 + 40, 1875 + 60, and 1875 + 80.
1935 m br	
1955 m br	
2010 m	1020 x 2, 1020 + 1219 and related peaks
2030 m	
2070 w	
2110 w	
2130 w	
2160 m	
2200 m	
2275 m sh	
2300 ms	
2350 ms	
2380 m sh	
2430 m	
2470 m	

TABLE 2 (cont'd)

2545 m	2 x 1267, 2 x 1312,
2590 m	1267 + 1312 and related
2620 m	peaks
2680 w	
2710 w	2 x 1592 and related peaks
2760 w	
2780 w	

---






Fig 1 Structure of the molecule at a crystal site; 'a' and 'e' indicate the axial and equatorial nitramino groups. A single line represents a single bond, a single line with dots a partial double bond, a heavy line a double bond




Fig 2 % Transmission (T) vs frequency traces of the infrared spectra of RDX in the 400 to 4000  $\text{cm}^{-1}$  region: (a) Spectrum of a thin single crystal (b) Spectrum of a thick single crystal in the 1600 to 4000  $\text{cm}^{-1}$  region (c) Spectrum of RDX in  $\text{d}_6$ -DMSO in the 1200 to 4000  $\text{cm}^{-1}$  region and RDX in acetonitrile in the 600 to 1200  $\text{cm}^{-1}$  region (d) Spectrum of KBr pellet of RDX




Fig 3 Traces of the Raman spectra of RDX in the 400 to 4000  $\text{cm}^{-1}$  region: (a) Raman spectrum in the 400 to 1500  $\text{cm}^{-1}$  region using 6328 Å excitation and slit width about 5  $\text{cm}^{-1}$  of a single crystal of RDX (b) Raman spectrum between 400 and 4000  $\text{cm}^{-1}$  of powdered RDX using 4880 Å excitation and slit width of about 5  $\text{cm}^{-1}$  (bands marked G are due to the glass capillary tube) (c) Raman spectrum between 400 and 1000  $\text{cm}^{-1}$  of  $\text{d}_6$ -DMSO solution of RDX using 4880 Å excitation and slit width of about 7  $\text{cm}^{-1}$  (bands marked S are due to the solvent)




Fig 4 % Transmission (T) vs frequency traces of the KBr pellet infrared spectra between 400 and 1700  $\text{cm}^{-1}$  of (a) Nitro group labeled RDX (b) Ring labeled RDX (c) Unlabeled RDX




Fig 5 Traces of the low frequency infrared and Raman spectra:  
 (a) Infrared spectra of pure RDX in Nujol mull, between 200 and 500 cm<sup>-1</sup> (b) Infrared spectra of the trinitroso analog of RDX, TNTC, in Nujol mull between 200 to 500 cm<sup>-1</sup> (c) Raman spectrum of an RDX single crystal using 6328 Å excitation between 30 and 400 cm<sup>-1</sup>. Slits of about 3 cm<sup>-1</sup> were used between 30 and 200, and of about 6 cm<sup>-1</sup> between 200 and 400 cm<sup>-1</sup> (d) Raman spectrum of powder RDX using 4830 Å radiation between 30 and 200 cm<sup>-1</sup>. Slits of about 3 cm<sup>-1</sup> were used.